

## Extreme Problem 7 Solutions

(i) Let the distance from the top surface of the cylindrical tank to the vertex of the cone structure

be  $x$ . Then  $\tan 60^\circ = \frac{r}{x} \Rightarrow x = r \cot 60^\circ = \frac{r}{\sqrt{3}}$

Height of cylindrical tank is therefore  $h - x = h - \frac{r}{\sqrt{3}}$  and

Volume of tank  $V_T = \pi r^2 \left( h - \frac{r}{\sqrt{3}} \right) = \pi r^2 h - \frac{1}{\sqrt{3}} \pi r^3$

When the cylindrical tank has a maximum volume,  $\frac{dV_T}{dr} = 0$  (Note:  $h$  is a constant)

$$2\pi r h - \sqrt{3} \pi r^2 = 0$$

$$\pi r (2h - \sqrt{3}r) = 0$$

$$\therefore r = \frac{2}{\sqrt{3}} h = \frac{2\sqrt{3}}{3} h \text{ (shown) or } r = 0 \text{ (NA)}$$

(ii) Let the slant height of the liquid chemical occupying the inverted cone at any time instant  $t$

be  $l$ . Since the semi-vertical angle of the said inverted cone is  $45^\circ$ , we have  $l = \sqrt{2}r$ .

Also, vertical height of liquid chemical in cone shall be equivalent to  $r$ .

As such, lateral surface area of cone occupied by liquid chemical at time  $t$  is given by

$$A = \pi r l = \sqrt{2}\pi r^2 \text{ -----(1)}$$

$$\text{Volume of cone } V = \frac{1}{3} \pi r^2 \times \text{height of liquid chemical} = \frac{1}{3} \pi r^3 \text{ -----(2)}$$

$$\text{From (1), } r = \left( \frac{A}{\sqrt{2}\pi} \right)^{\frac{1}{2}}; \text{ substituting this into (2) gives } V = \frac{1}{3} \pi \left( \frac{A}{\sqrt{2}\pi} \right)^{\frac{3}{2}} = \frac{2^{\frac{3}{4}}}{3\sqrt{\pi}} A^{\frac{3}{2}}$$

$$\text{Therefore, } \frac{dV}{dA} = \frac{3}{2} \cdot \frac{2^{\frac{3}{4}}}{3\sqrt{\pi}} A^{\frac{1}{2}} = \frac{2^{\frac{7}{4}}}{\sqrt{\pi}} A^{\frac{1}{2}}$$

After 30 minutes, volume of liquid chemical leaked into inverted cone =  $30 \times 0.3 = 9 \text{ m}^3$

$$\frac{1}{3} \pi r^3 = 9 \Rightarrow r = \left( \frac{27}{\pi} \right)^{\frac{1}{3}} \quad \text{and} \quad A = \sqrt{2\pi} \left( \frac{27}{\pi} \right)^{\frac{2}{3}} = 9\sqrt{2}\pi^{\frac{1}{3}} \text{ m}^2$$

$$\frac{dV}{dt} = \frac{dV}{dA} \times \frac{dA}{dt} \Rightarrow 0.3 = \frac{2^{\frac{7}{4}}}{\sqrt{\pi}} A^{\frac{1}{2}} \times \frac{dA}{dt}$$

$$\text{Since } A = 9\sqrt{2}\pi^{\frac{1}{3}} \text{ m}^2, \quad 0.3 = \frac{2^{\frac{7}{4}}}{\sqrt{\pi}} \left( 9\sqrt{2}\pi^{\frac{1}{3}} \right)^{\frac{1}{2}} \times \frac{dA}{dt}$$

$$= \frac{2^{\frac{7}{4}}}{\sqrt{\pi}} (3)(2)^{\frac{1}{4}} \pi^{\frac{1}{6}} \times \frac{dA}{dt}$$

$$= 3(2)^{\frac{3}{2}} \pi^{-\frac{1}{3}} \times \frac{dA}{dt}$$

$$\text{Hence, } \frac{dA}{dt} = \frac{3}{10} \div \frac{3}{2\sqrt{2}} \pi^{-\frac{1}{3}} = \frac{\sqrt{2}}{5} \pi^{\frac{1}{3}} \text{ m}^2 / \text{min (shown)}$$