

Binomial Series Summary

1. Expansion of $[1 + f(x)]^n$

This is given by

$$1 + nf(x) + \frac{(n)(n-1)}{2!}[f(x)]^2 + \frac{(n)(n-1)(n-2)}{3!}[f(x)]^3 + \frac{(n)(n-1)(n-2)(n-3)}{4!}[f(x)]^4 + \dots$$

Example: $(1 - 3x)^{-2} = 1 + (-3x)(-2) + \frac{(-2)(-3)}{2!}(-3x)^2 + \frac{(-2)(-3)(-4)}{3!}(-3x)^3 + \dots$

$$= 1 + 6x + 27x^2 + 108x^3 + \dots$$

Note:

(i) For $n \in \mathfrak{R}$, $n \neq \mathbb{Z}^+$, the structure intended for expansion **must strictly adhere to the form**

$[1 + f(x)]^n$. For instance, $(8 + 2x)^{\frac{1}{3}}$ must be reorganized as $8^{\frac{1}{3}}\left(1 + \frac{x}{4}\right)^{\frac{1}{3}}$ at the very beginning

before proceeding with the actual binomial series expansion.

(ii) The expansion of $[1 + f(x)]^n$ is **valid** for $|f(x)| < 1$.

2. Typical applications of the binomial series expansion

(a) To provide an estimation of specific surd values:

Example: $(8 - x)^{\frac{1}{3}} \approx 2 - \frac{1}{12}x + \frac{1}{288}x^2$; by substituting $x = \frac{1}{8}$, we can arrive at a value for $\sqrt[3]{63}$:

$$\left(\frac{63}{8}\right)^{\frac{1}{3}} = 2 - \frac{1}{12}\left(\frac{1}{8}\right) + \frac{1}{288}\left(\frac{1}{64}\right) = 2 - \frac{1}{96} + \frac{1}{18432}$$

$$\frac{1}{2}\left(\sqrt[3]{63}\right) = \frac{36673}{18432}$$

$$\text{Hence, } \sqrt[3]{63} = \frac{36673}{9216} \text{ (shown)}$$

(b) To compute the approximate equations of tangents to points on a sophisticated curve:

Example: if we are aware that $y = \left(\frac{1-x}{1+x}\right)^{\frac{1}{4}} \approx 1 - \frac{1}{2}x + \frac{1}{8}x^2$,

$$\text{Then } \frac{dy}{dx} \approx \frac{d}{dx} \left(1 - \frac{1}{2}x + \frac{1}{8}x^2\right) = -\frac{1}{2} + \frac{1}{4}x$$

Hence, at the point $\left(\frac{1}{3}, \frac{1}{\sqrt[4]{2}}\right)$, gradient of the tangent to the curve $y = \left(\frac{1-x}{1+x}\right)^{\frac{1}{4}}$ is approximately equal to $-\frac{1}{2} + \frac{1}{4}\left(\frac{1}{3}\right) = -\frac{5}{12}$, and the equation of this tangent is simply given by

$$y - \frac{1}{\sqrt[4]{2}} = -\frac{5}{12} \left(x - \frac{1}{3}\right), \text{ ie } y = -\frac{5}{12}x + \left(\frac{5}{36} + \frac{1}{\sqrt[4]{2}}\right) \text{ (shown)}$$

3. Finding the formula of the general term in the expansion

Example: In the expansion of $(x^2 + 4)^{-2} = \frac{1}{16} \left(1 + \frac{x^2}{4}\right)^{-2}$, coefficient of x^{2n}

$$= \left(\frac{1}{16}\right) \frac{(-2)(-3)(-4)\dots\dots(-n-1) \left(\frac{1}{4}\right)^n}{n!} = \left(\frac{1}{16}\right) \frac{(-1)^n (2)(3)(4)\dots\dots(n+1) \left(\frac{1}{4}\right)^n}{n!}$$

$$= \left(\frac{1}{16}\right) \frac{(-1)^n (n+1)! \left(\frac{1}{4}\right)^n}{n!} = \frac{(-1)^n (n+1)}{4^n (16)} = \frac{(-1)^n (n+1)}{4^{n+2}} \text{ (shown)}$$

4. Ascending series expansion

Example: $\left(1 + \frac{1}{4x^2}\right)^{\frac{1}{2}}$ may resemble the required format $[1 + f(x)]^n$, but if an ascending series is to

be achieved through expansion, then $f(x) = \frac{1}{4x^2}$ will not be appropriate as the main variable x resides purely in the denominator. A transformation has to be made:

$$\left(1 + \frac{1}{4x^2}\right)^{\frac{1}{2}} = \left(\frac{1}{4x^2}\right)^{\frac{1}{2}} (4x^2 + 1)^{\frac{1}{2}}$$

$$= (2x) \left[1 + \left(-\frac{1}{2}\right)(4x^2) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}(4x^2)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}(4x^2)^3 + \dots \right]$$

⋮